



E-ISSN: 2707-8353
P-ISSN: 2707-8345
IJCRO 2022; 4(2): 04-08
Received: 06-04-2022
Accepted: 10-05-2022

TJM Wetzels
MD, Department of
Orthopedic Surgery and
Traumatology, Sint-
Andriesziekenhuis Tielt,
Bruggestraat, Tielt, Belgium

The treatment of an Allen type III fingertip injury with a composite graft: A case report and review of the literature

TJM Wetzels

DOI: <https://doi.org/10.22271/27078345.2022.v4.i2a.114>

Abstract

Fingertip injuries are frequently seen and there are many treatment options. Small defects with limited exposure of the bone can be treated conservatively with healing by secondary nature. Good results can be achieved with standard dressings or a semi-occlusive dressing. With larger defects, surgical treatment is needed. Local advancement flaps can be used when there is sufficient local tissue. If this is not present, a pedicle flap or tissue transfer is needed. The non-microsurgical re-fixation of the amputated part as a composite graft yields excellent results in children. It is, however, not frequently used in adults, due to the less favourable results.

In this case report, an Allen type III fingertip amputation was successfully treated by non-microsurgical re-fixation of the amputated fingertip. At final follow up, 18 months after injury, there was a satisfactory aesthetic and functional result. Although smaller in size, the fingertip resembled the other fingers with normal nail growth. Sensibility had recovered moderately and cold intolerance was present. However, the patient was very satisfied with the result and did not experience any limitations during daily activities.

In conclusion, the use of a composite graft in Allen type III fingertip injuries is a viable option in selected cases.

Keywords: Fingertip injury, fingertip amputation, composite graft

Introduction

The finger is the most injured part of the upper extremity, accounting for more than a third of the injuries ^[1]. Although amputations of the finger only constitute a small portion of these injuries (2%), the burden of disability is highest, when compared to other injuries such as fractures ^[2]. The estimated incidence has been steady for the last decades with an average of 7, 5 injuries per 100.000 persons per year ^[1, 3]. The incidence rises with age in adults ^[3, 4]. Most non-occupational injuries occur at home due to a crushing or cutting trauma, such as a door or power tool ^[4]. Although the overall need for hospitalization is low, it is increasing ^[3, 4].

Distal amputations of the finger can be described using various classification systems. Most classification systems describe a transverse amputation level ^[5-8]. The Allen classification⁵ is based on the level of amputation with regard to the nail. In type I injuries there is only loss of the pulp. When there is loss of pulp and nail, it is classified as a type II. Type III injuries include partial loss of the terminal phalanx. And in type IV injuries the lunula of the nail is involved.

In the classification by Evans ^[9] a more extensive approach is used, classifying the injury to each individual component of the fingertip: pulp, nail and bone. This leads to a 3-digit code which provides detailed information about the injury. Using this classification system 648 combination can be made. Which reflects the great variation in injury patterns of fingertip amputations. As a result, there are great differences in the preferred treatment of fingertip amputations among international hand- and microsurgeons ^[10-12].

When primary closure is not possible, treatment options include: healing by secondary nature, revision amputation and local or regional flap coverage. The use of a composite graft has been proven to give good results in children ^[13, 14]. However, it is not frequently used in adults, due to the possible high failure rate ^[15].

I believe that composite grafting is a good alternative for treatment of Allen type III fingertips injuries in adults. Which is illustrated by the following case report and review of the literature.

Corresponding Author:
TJM Wetzels
MD, Department of
Orthopedic Surgery and
Traumatology, Sint-
Andriesziekenhuis Tielt,
Bruggestraat, Tielt, Belgium

Case Report

Our patient is a healthy 55-year-old man. He is a right-handed auto mechanic and a non-smoker. He injured the 5th finger of his left hand at home. He was chopping wood with the use of an automatic wood cleaver machine. This injury resulted in a fingertip amputation. Both his wife and daughter were trained nurses, they immediately put the amputated fingertip on ice (no direct contact). He presented to our emergency department within 1 hour of the injury.

Inspection of the finger showed a volar oblique amputation, starting just distal to the level of the lunula dorsally and ending distal to the DIP crease on the volar side. The distal phalanx was almost completely intact and protruded beyond the level of the wound. The amputated part consisted of the pulp, the majority of the nail bed and the complete nail (Figure 1). Due to the intact germinal matrix on the dorsal side, the injury was classified as an Allen type III.

Intravenous antibiotics were administered in the emergency department and sterile dressings were applied. Within 2 hours of presentation, the patient was prepared for surgery. After discussing the various treatment options, a non-microsurgical re-fixation was chosen.

Surgery was performed under loco-regional anaesthesia (supraclavicular plexus) and tourniquet control. The amputated part was cleaned and debrided with removal of the nail. Inspection showed a limited part of the tuft, which was removed. Then the proximal stump was first cleaned and debrided. Inspection showed a distal remnant of the ulnar digital nerve. It was resected deep to the level of the wound to prevent the formation of a neuroma. At this level no digital artery was seen. Due to the oblique nature of the wound, on the radial side no digital nerve or artery was found. Small incisions were made on the ulnar and radial corner of the nail fold to expose the proximal nail bed and germinal matrix. The amputated part was positioned on the exposed distal phalanx and the nail bed was sutured with absorbable 4/0 sutures (Vicryl Rapide, Ethicon Inc.). The skin on the radial, volar and ulnar side was sutured with a non-absorbable nylon 4/0 suture (Ethilon, Ethicon Inc.). A fake nail was made out of a 10cc syringe and held in place with a dorsal "figure of 8" nylon 2/0 suture (Figure 2). A sterile dressing was applied.

Postoperatively the patient was treated with intravenous antibiotics for 24 hours. After which he was discharged home with oral antibiotics for 1 week. Two days after discharge he was seen in our outpatient clinic and the dressings were removed. Some discoloration of the tip had appeared, without signs of infection. Daily wound care was started by his wife and daughter.

He was seen repeatedly in the outpatient clinic in the days and weeks after. 24 days after surgery the sutures and fake nail were removed (Figure 3). At this time the discoloration had progressed and the first signs of necrosis were visible at the volar and ulnar aspect. Due to stiffness of both the PIP and DIP joint, he was advised to start physiotherapy. During the next weeks the necrotic graft regressed in size, with proximal regeneration of the tissue (Figure 4). The graft was left in place to function as a biological dressing. After 4 months nail growth was observed and a two weeks later the remaining necrotic tip had fallen off. Functionally he did not experience any limitations with a good flexion and contact pulp-palm.

The patient was last seen at 18 months postoperatively (Figure 5). He was very satisfied with both the appearance and functional result. The length and mobility made it possible for him to perform his normal activities with excellent grip strength. He reported some loss of sensibility

and mild cold intolerance of the fingertip. On clinical examination there was a 2-point discrimination of 12mm. Range of motion of the MP and PIP joint equalled the contralateral side. There was only a slight decrease in flexion of the DIP joint, 52° compared to 60°. The length of the distal phalanx on the affected side was 1 cm shorter; 16mm versus 26mm (measured on the volar surface from the DIP crease).



Fig 1: (A) Intra-operative view of the injured finger. Notice the protruding bone. (B) Intra-operative view of the amputated fingertip with nail in situ




Fig 2: (A) Volar aspect after re-fixation of the fingertip. (B) Dorsal view with "fake" nail in situ



Fig 3: Volar view after suture removal at 3 weeks



Fig 4: Progression of healing / necrosis. (A) Volar view at 6 weeks. (B) Volar view at 8 weeks. (C) Volar view at 12 weeks. (D) Volar view at 18 weeks




Fig 5: Appearance of the injured finger compared to contralateral at 18 months. (A) Volar view; (B) Dorsal view

Types of treatment

Conservative

Conservative treatment of fingertip injuries provides good results with almost no difference in sensibility, range of motion or grip strength. Donor site morbidity and long immobilization are avoided. Which leads to an overall quick recovery, with an average healing time of 4 weeks and subsequent early return to work [5, 16, 17]. Treatment can easily be done at home, with for example the use of Vaseline and Coban tape. It is important to keep the wound bed moist [18].

Another form of healing by secondary nature is with the use of semi-occlusive dressings. Excellent results have been reported without complications or the need for secondary surgery [19]. Time to healing seems a little longer, averaging 6.5 weeks. However, it provides a high percentage of soft tissue regeneration of almost 90% [20]. Secondary healing seems to be the treatment option of choice in fingertip amputations, even with bone exposure up to the wound level. However, when the bone protrudes beyond the level of the wound, a surgical treatment seems necessary [19].

Surgical

When performing a revision amputation, the bone is shortened to the level where primary closure is possible. It provides good results when tension free closure is achieved with an average time to work of 7 weeks [21]. Sensibility is near to normal and range of motion is acceptable. However, cold intolerance is frequently seen with an average of 24%.²¹ Care must be taken if excessive shortening of the bone is required. A shorter phalanx increases the risk of nail deformity and can lead to serious functional deficits [22].

Therefore, numerous reconstruction methods with flap surgery have been described with the purpose of maintaining length and sensation. Various algorithms to achieve this goal have been described [23, 24]. They are based on the involved finger and geometry of the defect.

In small defects various types of VY advancement flaps can be performed. Volarly with the Atasoy flap, and laterally with the Kutler flap [25]. These flaps are not technically demanding and yield good results with limited complications. In medium sized defects other local advancement flaps can be used, depending on the geometry of the defect and affected finger. The Moberg flap, the Venkataswami flap and the Hueston flap are some examples [23, 26]. In a larger defect a pedicle flap is frequently needed.

The homodigital island flap, antegrade or retrograde, is frequently used in this situation [23]. The survival rate of these pedicle flaps is high, with normal sensibility and absence of cold intolerance [27, 28]. The first dorsal metacarpal artery (DMCA) or kite flap is another example

of a versatile pedicle flap [26]. Various other options include the cross-finger flap and the nar flap [23, 29]. Both are reliable and easy techniques.

Free tissue transfer with the use of a full thickness skin graft from the hypothenar region has also been described [30, 31]. However, it is commonly seen as a last resort, due to superior results of other techniques. Alternatively, a free pulp toe transfer can be performed. The transferred tissue resembles that of the fingertip, yielding good functional and esthetical results. With minimal donor site morbidity [29, 32].

Composite graft

Non-microsurgical refixation of the amputated fingertip is a common practice in children, although the clinical failure rate reported is high (up to 32% and 41%) [13, 14]. The patient reported survival, however, is much higher; 78.4% versus 59% [13]. And revision surgery is seldom needed because almost all of the failures heal by secondary nature [14]. Kiuchi *et al.* [33] reported similar results for the survival rate of the composite grafts in adults. They concluded that clean-cut injuries distal to the base of the nail (meaning Allen type I-III) are good candidates for composite grafting.

Discussion

In this case of an Allen type III fingertip amputation, the non-microsurgical re-fixation resulted in an excellent patient reported outcome. It is important to understand that the circumstances were optimal. It was a healthy, non-smoking patient. His relatives were trained nurses, who aided in the primary care of the amputated part and the precise aftercare. The surgery was also performed within several hours of the injury. If any of these factors had been different, the result might not have been as good. Clinical evaluation at 18 months revealed only moderate recovery of the sensibility, with the presence of cold intolerance. However, this was well tolerated by the patient, probably because it was the 5th finger. The regaining of his grip strength was more important to him than the loss of sensibility. A major drawback of the treatment was the long leave of absence from work (4.5 months). This was mainly a result of his job as an auto mechanic. The patient was very concerned that contamination with grease would have a negative impact on the healing.

From a clinical point of view, the graft survival can be classified as a failure. After 3 weeks the first signs of necrosis were visible and after 6 weeks almost the whole amputated portion had gone black. Although in the end more than 50% of the fingertip had regenerated with the presence of a nail. This result is probably achieved by a combination of graft take and secondary healing.

Various techniques have been described to enhance the outcome of the composite graft. These techniques include the use of cooling, hypobaric oxygen, and intravenous lipoprostaglandin E1 [34-37]. These techniques however, require a long hospital stay of up to 15 days. Microsurgical repair of an artery and vein also increases the survival rate [38, 39]. However, distal from the lunula (Allen type I-III) this reconstruction is described as very difficult to impossible [39].

Few other treatment options would have resulted in a better patient reported outcome in this case. With revision amputation the finger would have been significantly shortened, probably to the level of the DIP joint. Which would have probably resulted in loss of function [22]. A local

advancement flap was not possible due to the size and geometry of the defect. And although the sensibility might have been better with the use of a homo-digital island flap. The volar scar would have probably caused complaints with loss of mobility and grip strength ^[23]. An important advantage of the chosen treatment was the preservation of the nail. Which would not have been possible with the other mentioned treatment options.

In conclusion, a non-microsurgical re-fixation of the amputated fingertip as a composite graft in Allen type III injuries can be a viable treatment option in certain cases.

References

- Ootes D, Lambers KT, Ring DC. The epidemiology of upper extremity injuries presenting to the emergency department in the United States. *Hand (NY)*. 2012 Mar;7(1):18-22. doi: 10.1007/s11552-011-9383-z. Epub 2011 Dec 14. PMID: 23449400; PMCID: PMC3280373.
- Crowe CS, Massenburg BB, Morrison SD, Chang J, Friedrich JB, Abady GG, *et al*. Global trends of hand and wrist trauma: a systematic analysis of fracture and digit amputation using the Global Burden of Disease 2017 Study. *Inj Prev*. 2020 Oct;26(Suppl 1):i115-i124. doi: 10.1136/injuryprev-2019-043495. Epub 2020 Mar 13. PMID: 32169973; PMCID: PMC7571361.
- Reid D, Shah K, Eltorai A, Got C, Daniels A. Epidemiology of Finger Amputations in the United States From 1997 to 2016. *Journal of Hand Surgery Global Online*. April 2019;1(2):45-51. 10.1016/j.jhsg.2019.02.001.
- Conn JM, Annest JL, Ryan GW, Budnitz DS. Non-work-related finger amputations in the United States, 2001-2002. *Ann Emerg Med*. 2005 Jun;45(6):630-5. doi: 10.1016/j.annemergmed.2004.10.012. PMID: 15940097.
- Allen MJ. Conservative management of finger tip injuries in adults. *Hand*. 1980 Oct;12(3):257-65. doi: 10.1016/s0072-968x(80)80049-0. PMID: 7002744.
- Foucher G, Norris RW. Distal and very distal digital replantations. *Br J Plast Surg*. 1992 Apr;45(3):199-203. doi: 10.1016/0007-1226(92)90076-a. PMID: 1596659.
- Hirase Y. Salvage of fingertip amputated at nail level: new surgical principles and treatments. *Ann Plast Surg*. 1997 Feb;38(2):151-7. doi: 10.1097/00000637-199702000-00009. PMID: 9043584
- Tamai S. Twenty years' experience of limb replantation-review of 293 upper extremity replants. *J Hand Surg Am*. 1982 Nov;7(6):549-56. doi: 10.1016/s0363-5023(82)80100-7. PMID: 7175124
- Evans DM, Bernardis C. A new classification for fingertip injuries. *J Hand Surg Br*. 2000 Feb;25(1):58-60. doi: 10.1054/jhsb.1999.0305. Erratum in: *J Hand Surg [Br]* 2000 Aug;25(4):414. PMID: 10763726.
- Kwon SH, Lao WW, Hsu AT, Lee CH, Hsu CC, Huang JJ, *et al*. The Preferred Management of a Single-Digit Distal Phalanx Amputation. *J Reconstr Microsurg*. 2020 May;36(4):301-310. doi: 10.1055/s-0039-1701013. Epub 2020 Feb 5. PMID: 32023640.
- Shauver MJ, Nishizuka T, Hirata H, Chung KC. Traumatic Finger Amputation Treatment Preference among Hand Surgeons in the United States and Japan. *Plast Reconstr Surg*. 2016 Apr;137(4):1193-1202. doi: 10.1097/01.prs.0000481301.25977.80. PMID: 27018674; PMCID: PMC5079431.
- Nishizuka T, Shauver MJ, Zhong L, Chung KC, Hirata H. A Comparative Study of Attitudes Regarding Digit Replantation in the United States and Japan. *J Hand Surg Am*. 2015 Aug;40(8):1646-56. 1656.e1-3. doi: 10.1016/j.jhsa.2015.05.026. PMID: 26213200; PMCID: PMC4800816.
- Borrelli MR, Dupré S, Mediratta S, Bisquera A, Greig A. Composite Grafts for Pediatric Fingertip Amputations: A Retrospective Case Series of 100 Patients. *Plast Reconstr Surg Glob Open*. 2018 Jun;19;6(6):e1843. doi: 10.1097/GOX.0000000000001843. PMID: 30276062; PMCID: PMC6157946.
- Murphy AD, Keating CP, Penington A, McCombe D, Coombs CJ. Paediatric fingertip composite grafts: Do they all go black? *J Plast Reconstr Aesthet Surg*. 2017 Feb;70(2):173-177. doi: 10.1016/j.bjps.2016.11.002. Epub 2016 Nov 19. PMID: 28010933.
- Lemmon JA, Janis JE, Rohrich RJ. Soft-tissue injuries of the fingertip: methods of evaluation and treatment. An algorithmic approach. *Plast Reconstr Surg*. 2008 Sep;122(3):105e-117e. doi: 10.1097/PRS.0b013e3181823be0. PMID: 18766028.
- Krauss EM, Lalonde DH. Secondary healing of fingertip amputations: a review. *Hand (NY)*. 2014 Sep;9(3):282-8. doi: 10.1007/s11552-014-9663-5. PMID: 25191157; PMCID: PMC4152443.
- Chavez-Galvan CR, Martínez-Pérez R, Flores-Alvarez E, Martínez-Pérez A. Comparative Analysis of a Modified Secondary Healing Protocol for Fingertip Amputations and Non-microsurgical Reconstruction Techniques. *Rev Bras Ortop (Sao Paulo)*. 2021 Oct;25;57(1):108-112. doi: 10.1055/s-0041-1735944. PMID: 35198117; PMCID: PMC8856846.
- Lalonde DH, Bouhtiauy J. Secondary Healing of Fingertip Amputations: Simple Wound Care Advice for Patients. *Plast Reconstr Surg Glob Open*. 2022 Jan;18;10(1):e4020. doi: 10.1097/GOX.0000000000004020. PMID: 35070592; PMCID: PMC8769127.
- Quadlbauer S, Pezzei C, Jurkowitsch J, Beer T, Keuchel T, Hausner T, *et al*. Der Okklusionsverband zur Behandlung von Allen III and IV Fingerkuppenverletzungen als Alternative zu lokalen Lappenplastiken [The semi-occlusive dressing in treating Allen III and IV fingertip injuries as an alternative to local skin flaps]. *Unfallchirurg*. 2017 Nov;120(11):961-968. German. doi: 10.1007/s00113-016-0237-6. PMID: 27638553.
- Hoigné D, Hug U, Schürch M, Meoli M, von Wartburg U. Semi-occlusive dressing for the treatment of fingertip amputations with exposed bone: quantity and quality of soft-tissue regeneration. *J Hand Surg Eur*. 2014 Jun;39(5):505-9. doi: 10.1177/1753193413489639. Epub 2013 May 21. PMID: 23695151.
- Wang K, Sears ED, Shauver MJ, Chung KC. A systematic review of outcomes of revision amputation treatment for fingertip amputations. *Hand (N Y)*. 2013 Jun;8(2):139-45. doi: 10.1007/s11552-012-9487-0. PMID: 24426910; PMCID: PMC3653002.
- Champagne L, Hustedt JW, Walker R, Wiebelhaus J, Nystrom NA. Digital Tip Amputations from the Perspective of the Nail. *Adv Orthop*. 2016;2016:1967192. doi: 10.1155/2016/1967192. Epub

- 2016 Nov 13. PMID: 27957344; PMCID: PMC5124444.
23. Lemmon JA, Janis JE, Rohrich RJ. Soft-tissue injuries of the fingertip: methods of evaluation and treatment. An algorithmic approach. *Plast Reconstr Surg.* 2008 Sep;122(3):105e-117e. doi: 10.1097/PRS.0b013e3181823be0. PMID: 18766028.
 24. Das De S, Sebastin SJ. Soft Tissue Coverage of the Digits and Hand. *Hand Clin.* 2020 Feb;36(1):97-105. doi: 10.1016/j.hcl.2019.09.002. PMID: 31757352.
 25. Chakraborty SS, Kala PC, Sahu RK, Dixit PK, Katrolia D, Kotu S. Fingertip Amputation Reconstruction with VY Advancement Flap: Literature Review and Comparative Analysis of Atasoy and Kutler Flaps. *World J Plast Surg.* 2021 Sep;10(3):8-17. doi: 10.29252/wjps.10.3.8. PMID: 34912662; PMCID: PMC8662676.
 26. Germann G, Rudolf KD, Levin SL, Hrabowski M. Fingertip and Thumb Tip Wounds: Changing Algorithms for Sensation, Aesthetics, and Function. *J Hand Surg Am.* 2017 Apr;42(4):274-284. doi: 10.1016/j.jhsa.2017.01.022. PMID: 28372640.
 27. Varitimidis SE, Dailiana ZH, Zibis AH, Hantes M, Bargiotas K, Malizos KN. Restoration of function and sensitivity utilizing a homodigital neurovascular island flap after amputation injuries of the fingertip. *J Hand Surg Br.* 2005 Aug;30(4):338-42. doi: 10.1016/j.jhsb.2005.04.014. PMID: 15936132.
 28. Hamdi MF. Les lambeaux d'avancement en ilot pulpaire homodactyle des doigts longs. À propos de 32 cas [Advancement fingertip homodigital neurovascular island flaps of long fingers. About 32 cases]. *Chir Main.* 2011 Apr;30(2):105-9. French. doi: 10.1016/j.main.2011.02.001. Epub 2011 Mar 23. PMID: 21507699.
 29. Panattoni JB, De Ona IR, Ahmed MM. Reconstruction of fingertip injuries: surgical tips and avoiding complications. *J Hand Surg Am.* 2015 May;40(5):1016-24. doi: 10.1016/j.jhsa.2015.02.010. Epub 2015 Mar 29. PMID: 25823622.
 30. Martin C, González del Pino J. Controversies in the treatment of fingertip amputations. Conservative versus surgical reconstruction. *Clin Orthop Relat Res.* 1998 Aug;(353):63-73. doi: 10.1097/00003086-199808000-00008. PMID: 9728160.
 31. Kawaiiah A, Thakur M, Garg S, Kawasmi SH, Hassan A. Fingertip Injuries and Amputations: A Review of the Literature. *Cureus.* 2020 May 26;12(5):e8291. doi: 10.7759/cureus.8291. PMID: 32601565; PMCID: PMC7317129.
 32. Spyropoulou GA, Shih HS, Jeng SF. Free Pulp Transfer for Fingertip Reconstruction-The Algorithm for Complicated Allen Fingertip Defect. *Plast Reconstr Surg Glob Open.* 2016 Jan 7;3(12):e584. doi: 10.1097/GOX.0000000000000569. PMID: 26894009; PMCID: PMC4727693.
 33. Kiuchi T, Shimizu Y, Nagasao T, Ohnishi F, Minabe T, Kishi K. Composite grafting for distal digital amputation with respect to injury type and amputation level. *J Plast Surg Hand Surg.* 2015;49(4):224-8. doi: 10.3109/2000656X.2015.1020314. Epub 2015 Mar 6. PMID: 25746850.
 34. Hirasé Y. Postoperative cooling enhances composite graft survival in nasal-alar and fingertip reconstruction. *Br J Plast Surg.* 1993 Dec;46(8):707-11. doi: 10.1016/0007-1226(93)90204-o. PMID: 8298786.
 35. Idone F, Sisti A, Tassinari J, Nisi G. Cooling Composite Graft for Distal Finger Amputation: A Reliable Alternative to Microsurgery Implantation. *In vivo.* 2016 Jul-Aug;30(4):501-5. PMID: 27381615.
 36. Lee Y, Heo JW, Moon JS, Kim SW, Kim J. Effects of hyperbaric oxygen on graft survival outcomes in composite grafting for amputated fingertip injury. *Arch Plast Surg.* 2020 Sep;47(5):444-450. doi: 10.5999/aps.2020.00381. Epub 2020 Sep 15. PMID: 32971596; PMCID: PMC7520242.
 37. Eo S, Hur G, Cho S, Azari KK. Successful composite graft for fingertip amputations using ice-cooling and lipo-prostaglandin E1. *J Plast Reconstr Aesthet Surg.* 2009 Jun;62(6):764-70. doi: 10.1016/j.bjps.2007.09.064. Epub 2008 Apr 14. PMID: 18407819.
 38. Hattori Y, Doi K, Ikeda K, Abe Y, Dhawan V. Significance of venous anastomosis in fingertip replantation. *Plast Reconstr Surg.* 2003 Mar;111(3):1151-8. doi: 10.1097/01.PRS.0000046497.88549.E0. PMID: 12621185.
 39. Sebastin SJ, Chung KC. A systematic review of the outcomes of replantation of distal digital amputation. *Plast Reconstr Surg.* 2011 Sep;128(3):723-737. doi: 10.1097/PRS.0b013e318221dc83. PMID: 21572379; PMCID: PMC3163033.